

Proliferation of *Pseudo-nitzschia brasiliana* and *P. cf. pseudodelicatissima* (Bacillariophyceae) in the Estero Santa Cruz, northern Gulf of California, Mexico

Proliferación de *Pseudo-nitzschia brasiliana* y *P. cf. pseudodelicatissima* (Bacillariophyceae) en el Estero Santa Cruz, norte del Golfo de California, México

Ismael Gárate-Lizárraga¹, Carlos Antonio Poot-Delgado^{2,3}, Erick Ramsés Ramírez-Castillo⁴ and Manuel Humberto Páez-Hernández⁵

¹Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas, Departamento de Plancton y Ecología Marina, Avenida Instituto Politécnico Nacional s/n, Col. Playa Palo de Santa Rita, La Paz, Baja California Sur 23096, México. igarate@ipn.mx

²Centro de Estudios del Mar, Departamento de Recursos Acuáticos, Km. 1 Carretera Campeche-Hampolol s/n, Col. Palmas, Campeche, Campeche 24027, México

³Laboratorio de Botánica Marina y Planctología, Instituto de Ciencias Marinas y Pesquerías, Universidad Veracruzana, Calle Hidalgo Nº 617, Col. Río Jamapa, Boca del Río. Veracruz, C.P. 94290, México

⁴Comité de Sanidad Acuícola del Estado de Sonora, Ébanos Nº 58, Col. Fuentes del Mezquital, Hermosillo, Sonora, México

⁵Dirección General de Protección contra Riesgos Sanitarios del Estado de Sonora, Paseo del Canal y Comonfort, Edificio Sonora, Sonora, México

Abstract.- A moderate bloom of *Pseudo-nitzschia* species occurred during collection of phytoplankton in Estero Santa Cruz, the State of Sonora, Mexico, in August 2012. The abundance of *Pseudo-nitzschia* in 2 samples was 84×10^3 and 160×10^6 cells L^{-1} . Results show the presence of 2 species, *P. brasiliana* and *P. cf. pseudodelicatissima*, which were responsible for a microalgae bloom. This is the first proliferation of *P. brasiliana* in the Gulf of California. *P. cf. pseudodelicatissima* was present in higher densities. A short morphological description is provided, including data from scanning electronic microscopy.

Key words: Diatom blooms, *Pseudo-nitzschia brasiliana*, *P. cf. pseudodelicatissima*, oyster farming, Gulf of California, Mexico

INTRODUCTION

Microalgae blooms are common natural phenomena along the both coasts of the Gulf of California (Cortés-Altamirano & Núñez-Pasten 1992, Alonso-Rodríguez & Ochoa 2004, Martínez-López *et al.* 2006, 2008; Gárate-Lizárraga *et al.* 2007, 2009). These blooms include dinoflagellates, diatoms, raphidophytes, cyanobacteria, silicoflagellates and ciliates (Cortés-Altamirano & Núñez-Pasten 1992, Gárate-Lizárraga *et al.* 2002, Band-Schmidt *et al.* 2005, Gárate-Lizárraga & Muñetón-Gómez 2008, Gárate-Lizárraga & Muciño-Márquez 2012). Dinoflagellates and diatoms are responsible for most of the blooms in the Gulf of California (Gárate-Lizárraga *et al.* 2001, 2006, Alonso-Rodríguez & Ochoa 2004). Species of diatoms, such as *Chaetoceros curvisetus* Cleve, *C. debilis* Cleve, *C. radicans* F. Schütt, *C. socialis* H.S. Lauder, *Cylindrotheca closterium* (Ehrenberg) Reimann & J.C. Lewin, *Eucampia zodiacus* Ehrenberg, *Nitzschia sigma* (Kützing) W. Smith, *Pseudo-nitzschia* spp., *Rhizosolenia debyana* H. Peragallo, *Stephanopyxis palmeriana* (Greville) Grunow, and *Thalassiosira* spp.

have proliferated in the Gulf of California (Gárate-Lizárraga *et al.* 1990, 2001, 2003, 2006, Cortés-Altamirano & Núñez-Pasten 1992, Gárate-Lizárraga & Muñetón-Gómez 2009, Molina *et al.* 1997).

The potentially toxic diatom genus *Pseudo-nitzschia* is a common component of the microalgal assemblage in the Gulf of California (Gárate-Lizárraga *et al.* 1990, Moreno *et al.* 1996). *Pseudo-nitzschia* blooms in the study region are related to upwelling events (Gárate-Lizárraga *et al.* 2007). At present, 7 species of *Pseudo-nitzschia* have been identified in the Gulf of California: *P. americana* (Hasle) Fryxell, *P. australis* Frenguelli, *P. brasiliana* N. Lundholm, G.R. Hasle & G.A. Fryxell, *P. fraudulenta* (Cleve) Hasle, *P. pseudodelicatissima* (Hasle) Hasle, *P. pungens* (Grunow ex Cleve) Hasle, and *P. subfraudulenta* (G.R. Hasle) G.R. Hasle (Hernández-Becerril 1998, Lundholm *et al.* 2002, Sierra-Beltrán *et al.* 2005, Gárate-Lizárraga *et al.* 2007). Blooms of *Pseudo-nitzschia* have been scarcely reported in the Gulf of California (Sierra-

Beltrán *et al.* 1997, 2005, Cortés-Altamirano & Licea-Durán 2004, Gómez-Aguirre *et al.* 2004, Gárate-Lizárraga *et al.* 2007, Ayala-Rodríguez 2008). This report describes the first proliferation of the potentially toxic diatoms *Pseudo-nitzschia brasiliana* and *P. cf. pseudodelicatissima* in the Estero Santa Cruz, Mexico (August 2012), including a brief description of both taxa.

MATERIALS AND METHODS

The Estero Santa Cruz lies between the delta region of the Rio Colorado and the extensive mudflats of Ensenada Pabellones and Bahía Santa María, Sinaloa, approximately 1,000 km south of the Delta (Meling-López *et al.* 1998, Fleischner & Gates 2009). Estero Santa Cruz is a medium-sized lagoon south of the town of Bahía de Kino. Its 3,622 ha area contains salt marshes, mangroves, and sand/mud flats, as well as permanent channels up to 3 m in depth (Fleischner & Gates 2009). Since the damming of the Río Sonora in 1947, Estero Santa Cruz no longer receives freshwater. The estuary-turned-lagoon is hypersaline (Quevedo-Estrada 2007). This aquatic system is now developing into a region of aquaculture ponds for oyster farming and shrimp cultivation.

As part of a monitoring program, phytoplankton bottle samples were collected on 1 August 2012 at 2 sampling stations in the northern part of the lagoon (Station 1: 28°47'37.88"N, 111°54'43.22"W and Station 2: 28°48'0.35"N, 111°55'1.56"W) where oysters are cultivated (Fig. 1). Phytoplankton surface samples for species identification and counting were collected with

plastic flasks. Surface vertical tows were made with a hand net (50 cm in diameter, 20 µm mesh size). A portion of each tow was immediately fixed with Lugol's acid solution. Temperature, salinity, and pH were recorded with a calibrated multi-parameter sampler (HI9828) at each sampling station. Cells were counted in 10 mL sedimentation chambers under an inverted microscope (Axio Observer, Carl Zeiss, Oberkochen, Germany; Hasle 1978). Primary identification was made up to the genus level and further some specimens were examined by scanning electron microscopy (JEOL JSM-5600 SEM, Peabody, Mass.). Samples were prepared for SEM as described in Kaczmarek *et al.* (2005). Preparation and observation was done at the Digital Microscopy Facility, Mount Allison University, Canada. In each sample, specimens identified as *Pseudo-nitzschia* spp. were examined for morphometric characteristics (width and length of the valve, density of striae, fibulae and poroids under SEM).

RESULTS AND DISCUSSION

Sea surface temperature during the bloom varied from 32.5 to 34°C. Salinity ranged from 34 to 37, and the pH ranged from 7.6-7.7. The species list and abundances are compiled in Table 1. Total phytoplankton abundance at Station 1 was 556×10^3 cells L⁻¹ and 618×10^3 cells L⁻¹ at Station 2. Nanoflagellates were numerically the most important group, followed by diatoms and dinoflagellates (Table 1). Among diatoms, abundance of *Pseudo-nitzschia* in the 2 samples was 84×10^3 at Station 2 and




Figure 1. Location of Estero Santa Cruz (~28°48'N, 111°55'W), and sampling sites adjacent to Bahía Kino, Sonora, Mexico in August 2012 / Localización del Estero Santa Cruz (~28°48'N, 111°55'W), y los sitios de muestreos adyacentes a Bahía Kino, Sonora, México en agosto de 2012

Table 1. Abundance of microalgae species recorded in the Estero Santa Cruz, Bahía Kino, Sonora during the proliferation of *Pseudo-nitzschia* spp. in August 2012 / Abundancia de las especies de microalgas registradas en el Estero Santa Cruz, Bahía Kino, Sonora durante la proliferación de *Pseudo-nitzschia* spp. en agosto de 2012.

Species composition	Sampling station (cells L ⁻¹)	
	1	2
Diatoms		
<i>Bacteriastrum</i> sp.	100	100
<i>Chaetoceros</i> spp.	29300	30700
<i>Chaetoceros peruvianus</i> Brightwell	1600	1000
<i>Cylindrotheca closterium</i> (Ehrenberg) Reimann & J.C.Lewin	22700	27000
<i>Eucampia zodiacus</i> Ehrenberg	100	0
<i>Guinardia striata</i> (Stolterfoth) Hasle in Hasle & Syvertsen	6000	8200
<i>Pseudo-nitzschia brasiliana</i> N. Lundholm, G.R.Hasle & G.A.Fryxell	68200	33700
<i>Pseudo-nitzschia</i> cf. <i>pseudodelicatissima</i> (Hasle) Hasle	92700	50400
<i>Navicula</i> spp.	600	0
<i>Rhizolenia setigera</i> Brightwell	27300	16000
<i>Rhizolenia</i> spp.	500	600
Centric unidentified diatoms	3600	4700
Pennate unidentified diatoms	6200	8600
Total abundance of diatoms	258900	181000
Dinoflagellates		
<i>Tripes furca</i> (Ehrenberg) F. Gómez,	0	100
<i>Prorocentrum micans</i> Ehrenberg	200	100
<i>Prorocentrum minimum</i> (Pavillard) J.Schiller	100	0
<i>Peridinium quinquecorne</i> Abé	0	300
<i>Protoperidinium</i> spp.	400	300
Naked dinoflagellates >20 µm	700	0
Naked dinoflagellates <20 µm	1100	1600
Total abundance of dinoflagellates	2500	2400
Nanoflagellates		
Total phytoplankton abundance	295200	435000
	556600	618400

160 x 10³ cells L⁻¹ at Station 1, respectively. Scanning electronic microscopy showed 2 species: *P. brasiliana*, with abundances varying in the 2 sampling stations from 33 x 10³ to 68 x 10³ cells L⁻¹, respectively, and *P. cf. pseudodelicatissima*, with abundances varying in the 2 sampling stations from 50 × 10³ to 92 × 10³ cells L⁻¹, respectively.

Pseudo-nitzschia brasiliana N. Lundholm, G.R. Hasle & G.A. Fryxell. SEM examination showed that this species has linear frustules in girdle view, rectangular shape in valve view, with widely rounded ends (Fig. 2 a-b) and lacks a central nodule (Fig. 2 c). *P. brasiliana* was mainly observed as single cells or forming chains of 2 or 3 cells. Morphological measurements are: 36.4-40.1 µm long, 2.07-

2.25 µm wide; 23-26 striae in 10 µm, and 22-26 fibulae in 10 µm, 7-9 poroids in 1 µm. The species morphometric data coincided with the original description and literature (Lundholm *et al.* 2002, Quijano-Scheggia *et al.* 2011, Sahraoui *et al.* 2011, and Parsons *et al.* 2012). *P. brasiliana* was first described in Brazil by Lundholm *et al.* (2002) and has also been reported in other warm water regions, such as the Gulf of Panama, Gulf of Mexico, central Mexican coast along the Pacific; Brazil, Indonesia, Malaysia, Thailand, South Korea, Tunisia, and Vietnam (Lundholm *et al.* 2002, Villac *et al.* 2005, Lim *et al.* 2010, Quijano-Scheggia *et al.* 2011, Sahraoui *et al.* 2011).

Pseudo-nitzschia cf. *pseudodelicatissima* (Hasle) Hasle was also identified. This species was found in both




Figure 2. Scanning electron micrographs of *Pseudo-nitzschia* spp. collected in the Estero Santa Cruz, Bahía Kino, State of Sonora, Mexico. (a-b) valves of *P. brasiliana*, scale bar = 5 μm ; (c) central part of the valve of *P. brasiliana* showing absence of central interspace, scale bar = 1 μm ; (d-e) valves of *P. cf. pseudodelicatissima*, scale bar = 10 μm ; (f) *P. cf. pseudodelicatissima* central interspace, scale bar = 1 μm / Imágenes de microscopio electrónico de *Pseudo-nitzschia* spp. colectadas en el Estero Santa Cruz, Bahía Kino, Sonora, México. (a-b) *P. brasiliana*; escala = 5 μm ; (c) parte central de la valva de *P. brasiliana* mostrando la ausencia de interspacio central, escala de la barra = 1 μm ; (d-e) *P. cf. pseudodelicatissima*, escala de la barra = 10 μm ; (f) *P. cf. pseudodelicatissima* mostrando el interspacio central, escala = 1 μm .

samples, but it was more abundant in sample 1 (Table 1). Examination by SEM showed that this species has completely linear and symmetrical valves, terminating at rounded apices (Figs. 2 d-e). *P. cf. pseudodelicatissima* was observed as single cells or forming chains of 2 or 3 cells. A central, larger interspace, corresponding to 4-6

striae, was observed (Fig. 2 f). *P. cf. pseudodelicatissima* cells are 55.6-82.3 μm long, 1.40-1.61 μm wide; 41-42 striae in 10 μm , 6-7 poroids in 1 μm , and 15-18 fibulae in 10 μm . Cells are linear in valve view, tapering part near the tips very short. Their morphometric characteristics and measurements are consistent and within the range of *P.*

pseudodelicatissima (Hasle & Syvertsen, 1997, Lundholm *et al.* 2002, Hong-Chang *et al.* 2012, Parsons *et al.* 2012).

At least 4 different species have been described in the literature in the *P. pseudodelicatissima/cuspidata* complex: *P. pseudodelicatissima*, *P. cuspidata* (Hasle) Hasle, *P. calliantha* Lundholm, Moestrup & Hasle, and *P. caciantha* Lundholm, Moestrup & Hasle (Lundholm *et al.* 2003). The *pseudodelicatissima/cuspidata* complex is distinguished by differences in the structure of the poroid hymens and girdle bands (Lundholm *et al.* 2003), separating *P. pseudodelicatissima* and *P. cuspidata* from *P. calliantha*. The hymen in *P. cuspidata* and *P. pseudodelicatissima* is similar and could be distinguished between these 2 species using transmission electron microscopy. Although *P. pseudodelicatissima* resembles *P. cuspidata*, our specimens are most likely not *P. cuspidata* because the valves are clearly linear and not lanceolate. They are distinguished by cell width, which is relatively thin in *P. pseudodelicatissima* (Lundholm *et al.* 2003).

In previous reports, toxicity of *P. brasiliana* in several strains from Brazil, Spain, and Malaysia were negative (Lundholm *et al.* 2003, Quijano-Scheggia *et al.* 2010); however, *P. brasiliana* from Bizerte Lagoon, Tunisia produces domoic acid (Sahraoui *et al.* 2011). *Pseudo-nitzschia pseudodelicatissima* can produce domoic acid, a causative agent of amnesic shellfish poisoning, notably some strains isolated from the Gulf of Mexico and Greece (Pan *et al.* 2001, Moschandreu *et al.* 2010). However, domoic acid has not been detected in strains of *P. pseudodelicatissima* from Gafahna, Portugal, Denmark Strait (northwest of Iceland), and Bizerte Lagoon, Tunisia (Lundholm *et al.* 2003, Sahraoui *et al.* 2011).

Given the occurrence of the potentially toxigenic diatoms *P. brasiliana* and *P. cf. pseudodelicatissima* at moderate cell densities in the Estero Santa Cruz, it appears that toxic events may occur in the future. Here we report the first proliferation of *P. brasiliana* in the Gulf of California. The first bloom of *P. pseudodelicatissima* was reported in Mazatlán, Sinaloa (8 July 2004). The death of ten brown pelicans (*Pelecanus occidentalis californicus* Ridgway), as well as several fish species were partly attributed to the *Pseudo-nitzschia* bloom (Sierra-Beltrán *et al.* 2005). In this study, we did not isolate either species of *Pseudo-nitzschia* to perform toxicity tests. A bloom of the 2 potentially toxic species of *Pseudo-nitzschia* in this lagoon system warrants a systematic and continuous

monitoring program by health authorities, because commercial shellfish aquaculture sites are located there.

ACKNOWLEDGMENTS

We thank the Sistema Producto Ostión en Sonora, and the Instituto Politécnico Nacional (Project: SIP-20121153) for their logistical and financial support for analysis and samples collection. We gratefully thank James Ehrman (Digital Microscopy Facility, Mount Allison University, Sackville, New Brunswick, Canada) for producing the scanning electron micrographs of *Pseudo-nitzschia* species used in this study. We thank M. C. Ramírez-Jáuregui (Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México) for helping us with the literature search. The authors thank the two anonymous reviewers for their helpful comments and suggestions.

LITERATURE CITED

- Alonso-Rodríguez R & JL Ochoa. 2004. Hydrology of winter-spring 'red-tides' in Bahía de Mazatlán, Sinaloa, Mexico. *Harmful Algae* 3: 163-171.
- Ayala-Rodríguez AG. 2008. Grupos funcionales del fitoplancton y estado trófico del Sistema Lagunar Topoloampo-Ohuira-Santa María. Tesis de Maestría. CICIMAR-IPN, La Paz, B.C.S., 149 pp.
- Band-Schmidt CJ, A Martínez-López & I Gárate-Lizárraga. 2005. First record of *Chattonella marina* in Bahía de La Paz, Gulf of California. *Harmful Algae News* 28: 6-7.
- Cortés-Altamirano R & S Licea-Durán. 2004. Decoloración en proliferaciones de microalgas como parámetro bioindicador en la Bahía de Mazatlán, México. *Revista de Biología Tropical* 52(Supl. 1): 27-34.
- Cortés-Altamirano R & A Núñez-Pasten. 1992. Doce años (1979-1990) de registros de mareas rojas en la Bahía de Mazatlán, Sinaloa, México. *Anales del Instituto de Ciencias del Mar y Limnología, UNAM* 19: 113-121.
- Fleischner TL & RH Gates. 2009. Shorebird use of Estero Santa Cruz, Sonora, Mexico: Abundance, diversity, and conservation implications. *Waterbirds* 32: 36-43.
- Gárate-Lizárraga I & RE Muciño-Márquez. 2012. Blooms of *Trichodesmium erythraeum* and *T. thiebautii* (Cyanobacteria, Oscillatoriales) in the Bahía de La Paz, Gulf of California. *CICIMAR, Oceanides* 27: 61-64.
- Gárate-Lizárraga I & MS Muñetón-Gómez. 2008. Bloom of *Peridinium quinquecorne* in the Ensenada de la Paz, Gulf of California (July 2003). *Acta Botanica Mexicana* 83: 33-47.

- Gárate-Lizárraga I & MS Muñetón-Gómez. 2009.** Primer registro de la diatomea epibionte *Pseudohimantidium pacificum* y de otras asociaciones simbióticas en el Golfo de California. *Acta Botanica Mexicana* 88: 33-47.
- Gárate-Lizárraga I, DA Siqueiros-Beltrones & CA Lechuga-Devéze. 1990.** Estructura de las asociaciones microfitoroplanctónicas de la región central del Golfo de California en el otoño de 1986. *Ciencias Marinas* 16: 131-153.
- Gárate-Lizárraga I, ML Hernández-Orozco, CJ Band-Schmidt & G Serrano-Casillas. 2001.** Red tides along the coasts of Baja California Sur, Mexico (1984-2001). *Oceanides* 16: 127-134.
- Gárate-Lizárraga I, C Band-Schmidt, R Cervantes-Duarte & DG Escobedo-Urías. 2002.** Mareas rojas de *Mesodinium rubrum* (Lohmann) Hamburger y Budenbrock ocurridas en el Golfo de California durante el invierno de 1998. *Hidrobiológica* 12: 15-20.
- Gárate-Lizárraga I, DA Siqueiros-Beltrones & V Maldonado-López. 2003.** First record of a *Rhizosolenia debyana* bloom in the Gulf of California, México. *Pacific Science* 57: 141-145.
- Gárate-Lizárraga I, MS Muñetón-Gómez & V Maldonado-López. 2006.** Florecimiento del dinoflagelado *Gonyaulax polygramma* frente a la isla Espíritu Santo, Golfo de California, México. *Revista de Investigaciones Marinas* 27: 31-39.
- Gárate-Lizárraga I, CJ Band-Schmidt, DJ López-Cortés & JJ Bustillos-Guzmán. 2007.** Bloom of *Pseudo-nitzschia fraudulenta* in Bahía de La Paz, Gulf of California (June-July 2006). *Harmful Algae News* 33: 6-7.
- Gárate-Lizárraga I, CJ Band-Schmidt, F Aguirre-Bahena & T Grayeb-del Álamo. 2009.** A multi-species microalgae bloom in Bahía de La Paz, Gulf of California, Mexico (June 2008). *CICIMAR, Oceanides* 24: 1-15.
- Gómez-Aguirre S, S Licea-Durán & S Gómez. 2004.** Proliferaciones de *Pseudo-nitzschia* spp. (Bacillariophyceae) y otras especies del microplancton en la Bahía de Mazatlán, México. *Revista de Biología Tropical* 52(Supl. 1): 69-76.
- Hasle GR. 1978.** The inverted-microscope method. In: Sournia A (ed). *Phytoplankton manual*, pp. 88-96. UNESCO, Paris.
- Hasle GR & EE Syvertsen. 1997.** Marine diatoms. In: Tomas CR (ed). *Identifying marine phytoplankton* pp. 1-385. Academic Press, San Diego.
- Hernández-Becerril DU. 1998.** Species of the planktonic diatom genus *Pseudo-nitzschia* of the Pacific coast of Mexico. *Hydrobiologia* 379: 77-84.
- Hong-Chang L, L Po-Teen, SS Nyun-Pau, Y Kotaki & L Chui-Pin. 2012.** Morphological observation of two species of *Pseudo-nitzschia* (Bacillariophyceae) Coastal Marine Science 35: 52-57.
- Kaczmarek I, JL Martin, MM LeGresley & JM Ehrman. 2005.** Diversity of the diatom genus *Pseudo-nitzschia* Peragallo in the Quoddy Region of the Bay of Fundy, Canada. *Harmful Algae* 4: 1-19
- Lim HC, SNP Su, MA Hartina, Y Kotaki, CP Leaw & PT Lim. 2010.** Toxicity of diatom *Pseudo-nitzschia* (Bacillariophyceae) analyzed using high performance liquid chromatography (HPLC). *Journal of Science and Technology in the Tropics* 6: 116-119.
- Lundholm N, GR Hasle, GA Fryxell & PE Hargraves. 2002.** Morphology, phylogeny and taxonomy of species within the *Pseudo-nitzschia americana* complex (Bacillariophyceae) with descriptions of two new species, *Pseudo-nitzschia brasiliensis* and *Pseudo-nitzschia linea*. *Phycologia* 41: 480-497.
- Lundholm N, O Moestrup, GR Hasle & K Hoef-Emden. 2003.** A study of the *Pseudo-nitzschia pseudodelicatissima/cuspidata* complex (Bacillariophyceae): what is *P. pseudodelicatissima*? *Journal of Phycology* 39: 797-813.
- Martínez-López A, CJ Band-Schmidt, D Escobedo-Urías & AE Ulloa-Pérez. 2006.** Bloom of *Chattonella subsalsa* in an impacted coastal lagoon in the Gulf of California. *Harmful Algae News* 31: 4-5.
- Martínez-López A, DC Escobedo-Urías, AE Ulloa-Pérez & R Aguirre. 2008.** Dynamics of a *Prorocentrum minimum* bloom along the northern coast of Sinaloa, México. *Continental Shelf Research* 28: 1693-1701.
- Meling-López AE, IA Villalba-Atondo & P Ortega-Romero. 1998.** Las halófitas de la Laguna Costera La Cruz. *Revista de Conservación de Suelo y Agua* 1: 1-7.
- Molina RE, FA Manrique & J García. 1997.** Nota sobre un florecimiento de *Stephanopyxis palmeriana* (Greville) Grunow (Bacillariophyceae) en la bahía Kun Kaak. *Golfo de California. Hidrobiológica* 7: 84-86.
- Moreno JL, S Licea-Durán & H Santoyo. 1996.** Diatomeas del Golfo de California, 273 pp. Universidad Autónoma de Baja California Sur-SEPFOMES-PROMARCO, México.
- Moschandreou KK, D Papaefthimiou, P Katikou, E Kalopesa, A Panou & G Nikolaidis. 2010.** Morphology, phylogeny and toxin analysis of *Pseudo-nitzschia pseudodelicatissima* (Bacillariophyceae) isolated from the Thermaikos Gulf, Greece. *Phycologia* 49: 260-273.
- Pan Y, ML Parsons, M Busman, P Moeller, Q Dortch, CL Powell, GA Fryxell & GJ Doucette. 2001.** *Pseudo-nitzschia pseudodelicatissima* - a confirmed producer of domoic acid from the northern Gulf of Mexico. *Marine Ecology Progress Series* 220: 83-92.
- Parsons ML, YB Okolodkov & JA Aké-Castillo. 2012.** Diversity and morphology of the species of *Pseudo-nitzschia* (Bacillariophyta) of the National Park Sistema Arrecifal Veracruzano, SW Gulf of Mexico. *Acta Botanica Mexicana* 98: 51-72.

- Quevedo-Estrada JL. 2007.** Propuesta de acción para la gestión integrada en la costa de Hermosillo, Sonora. Tesis de Especialidad en Gestión Integrada de Cuencas Hidrológicas. Colegio de Sonora, Hermosillo, 42 pp. <<http://www.colson.edu.mx/Cuencas/Documents/TESINA-JLQE.pdf>>
- Quijano-Scheggia S, E Garcés, K Andree, P de la Iglesia, J Diogène, JM Fortuño & J Camp. 2010.** *Pseudo-nitzschia* species in the Catalan Coast: characterization and contribution to the current knowledge of genus distribution in the Mediterranean Sea. *Scientia Marina* 74: 395-410.
- Quijano-Scheggia S, A Olivos-Ortiz, JH Gaviño-Rodríguez, F Castro-Ochoa, M Rivera-Vilarelle, MA Galicia-Pérez & M Patiño Barragán. 2011.** The first report of *Pseudo-nitzschia brasiliiana* and *P. micropora* (Bacillariophyceae) found in Cuyutlan Lagoon, Mexico. *Revista de Biología Marina y Oceanografía* 46: 189-197.
- Sahraoui I, SS Bates, D Bouchouicha, HH Mabrouk & AS Hlaili. 2011.** Toxicity of *Pseudo-nitzschia* populations from Bizerte Lagoon, Tunisia, southwest Mediterranean, and first report of domoic acid production by *P. brasiliiana*. *Diatom Research* 26: 293-303.
- Sierra-Beltrán AP, M Palafox-Uribe, J Grajales-Montiel, A Cruz-Villacorta & JL Ochoa. 1997.** Sea bird mortality at Cabo San Lucas, Mexico: evidence that toxic diatom blooms are spreading. *Toxicon* 35: 447-453.
- Sierra-Beltrán AP, R Cortés-Altamirano, JP Gallo-Reynoso, S Licea-Durán & J Égido-Villarreal. 2005.** Is *Pseudo-nitzschia pseudodelicatissima* toxin the principal cause of sardines, dolphins, sea lions and pelicans mortality in 2004 in Mexico? *Harmful Algae News* 29: 6-8.
- Villac MC, S Melo, M Menezes & D Rivera. 2005.** *Pseudonitzschia brasiliiana* (Bacillariophyceae), an opportunistic diatom on the coast of the state of Rio de Janeiro, Brazil. *Atlântica, Rio Grande* 27: 139-145.

Received 5 December 2012 and accepted 10 June 2013

Associate Editor: Pilar Muñoz M.