

# Length-weight relationship of four batoid species from the Pacific coast of Ecuador

Relación longitud-peso de cuatro especies de batoideos de la costa del Pacífico de Ecuador

Jonathan Pincay-Espinoza<sup>1,2\*</sup>, Fernando Rey Diz<sup>3</sup> and José Vélez-Tacuri<sup>4,5</sup>

<sup>1</sup>Maestría de Investigación en Acuicultura, Instituto de Posgrado, Universidad Técnica de Manabí, Bahía de Caráquez, Sucre, Manabí, Ecuador

<sup>2</sup>Subsecretaría de Recursos Pesqueros, Viceministerio de Acuicultura y Pesca, Puerto Pesquero Artesanal de San Mateo, Manta, Manabí, Ecuador

<sup>3</sup>World Wildlife Fund, Avenida Orellana E11-28 y Av. Coruña, Quito, Pichincha, Ecuador

<sup>4</sup>Fundación Red de Agentes por la Conservación y Sostenibilidad de los Ecosistemas, RACSE, Manta, Manabí, Ecuador

<sup>5</sup>Facultad Ciencias del Mar, Universidad Laica "Eloy Alfaro" de Manabí, Ciudadela Universitaria vía San Mateo, Manta, Manabí, Ecuador

\*Corresponding author: [jona.eduar@hotmail.com](mailto:jona.eduar@hotmail.com)

**Abstract.** Length-weight relationships (LWR) were estimated for four species of batoids, *Zapteryx xyster*, *Rostroraja equatorialis*, *Pseudobatos leucorhynchus* and *Gymnura crebripunctata*, collected in Ecuadorian waters. Values of the allometric growth parameter (*b*) oscillated from 2.41 to 3.15. *Pseudobatos leucorhynchus* and *G. crebripunctata* exhibited isometric type growth, while *Z. xyster* and *R. equatorialis* showed negative and positive allometric growth, respectively. New records of minimum sizes were obtained for free-swimming *R. equatorialis* (7.8 cm TL) and *G. crebripunctata* (20.0 cm DW). Data obtained constitute a baseline for future management and conservation studies.

**Key words:** Length-weight, batoids, Ecuadorian waters

## INTRODUCTION


Length-weight relationship (LWR) is a useful approach that provides basic data on population dynamics of the species, which facilitates their management and conservation through the formulation of fisheries management measures (Froese 2006, Koushlesh *et al.* 2017, Correia *et al.* 2018, Wei *et al.* 2019). LWR analysis is used to assess weight of an organism from a determined length (Froese 2006); understanding changes in fish weight as a function of size is essential information to deduce age structure, calculate growth rates or quantify some other aspect of fish population dynamics (Jellyman *et al.* 2013, Wang *et al.* 2013, Liu *et al.* 2018). Recently, a number of LWR of batoids have been reported (Romero-Caicedo *et al.* 2015, Rastgoo *et al.* 2016, Ehemann *et al.* 2017, Gladston *et al.* 2018, González-González & Ehemann 2019); however, life history of many batoids are still unknown or have been little studied (Dulvy *et al.* 2014).

In Ecuador, batoids are caught as part of bycatch of fishing fleets that operate with shrimp trawls and bottom gill nets. Most of these organisms are discarded and some are destined for local consumption or converted into fishmeal

(Martínez-Ortiz & García-Domínguez 2013). Information of this biological group is scarce. Thus, the current work aim to determine LWR parameters of four batoid species caught in Ecuadorian waters, being one of the first reports about life cycle of *Gymnura crebripunctata* (Peters, 1869) and *Rostroraja equatorialis* (Jordan & Bollman, 1890), and an increase in the biological knowledge of *Pseudobatos leucorhynchus* (Günther, 1867) and *Zapteryx xyster* Jordan & Evermann, 1896. This information is important for subsequent population assessment, allowing the proposal of management measures for these resources.

## MATERIALS AND METHODS

Specimens of the four batoid species were collected from shrimp trawlers as bycatch along the Ecuadorian coast (1°36'3"N and 79°12'2"W to 03°0'19"S and 80°44'4"W), between 20 and 80 m depth, from October 2016 through August 2017. A total of 12 fishing trips were monitored, information on four batoid species was registered randomly of 49 shrimp trawl spears. Trawl nets had a dimension between 30 and 35 m long and a mesh size of 3.8-5.0 cm. Trawl lasted between 3 and 4 h, with speeds of 1.8 to 2 kt.



All specimens were identified using the identification guide of Last *et al.* (2016). Sex was identified by the presence (males) or absence (females) of claspers. Size was recorded by means of a measuring tape with a precision of 0.1 cm; for each specimen, the total length (TL,  $\pm 0.1$  cm) was recorded from tip of rostral to distal point of caudal; in the case of *Gymnura crebripunctata*, the disc width (DW,  $\pm 0.1$  cm) was registered measured as the straight-line distance between the two outermost points of pectoral fins (Romero-Caicedo *et al.* 2015). Total weight (TW) was measured to the nearest gram (g) using a digital balance ( $W$ ,  $\pm 0.1$  g).

Total length was used for the length-weight relationship (LWR), except for *G. crebripunctata* where it was replaced by DW. LWR was described using a power model  $W = a \cdot TL^b$  or  $W = a \cdot DW^b$ , where  $a$  and  $b$  are parameters of the model estimated as the linear regression between  $\log(W)$  and  $\log(TL)$  or  $\log(DW)$ , meaning  $\log W = \log a + b \cdot \log TL$  or  $\log W = \log a + b \cdot \log DW$  (Froese 2006). 95% confidence limits were calculated for parameters  $a$  and  $b$ , as well as coefficient of determination ( $r^2$ ) to quantify variation of model explained by length-weight relationship. LWR was evaluated for males and females of each species and sexual differences were tested by a covariance analysis (ANCOVA), using size as a covariant (Zar 1999), with a significance level of 0.05. Type of growth for all species was determined by the slope  $b$ , when  $b = 3$ , the species presents an isometric growth, whereas allometric growth is negative or positive when  $b$  is  $< 3$  or  $> 3$ , respectively (Froese 2006), and it was evaluated using a Student's  $t$  test for slopes ( $H_0: b = 3$ ). All analyses were done using R (R Core Team 2013) and the package FSA (Ogle *et al.* 2018).

## RESULTS AND DISCUSSION

A total of 1,424 organisms of all species were analyzed. Results of the model for species and sex, sample size, minimum and maximum value of length and weight, estimates of parameters  $a$  and  $b$  with their confidence intervals ( $CI_{95\%}$ ), coefficient of determination ( $r^2$ ), Student's  $t$  test for slopes with their level of significance, and growth type are shown in Table 1.

Minimum total lengths herein reported were the smallest values for free-living organisms of *R. equatorialis* and *G. crebripunctata*, with 7.8 cm TL and 20.0 cm DW, respectively. Other studies carried out in the Eastern Pacific coasts record minimum sizes of 25.0 cm TL for *R. equatorialis* (Martínez-Ortiz & García-Domínguez 2013) and 29.0 cm DW for *G. crebripunctata* (Smith *et al.* 2009). The current study provides new minimum lengths for these species along the Ecuadorian coast; these new registers can be a result of fishing effort, combining two factors: catches in areas near

the coast (20-80 m of depth) and a small mesh size (3.8-5.0 cm), as smaller organisms can be entangled in fishing gears. Thus, elasmobranchs, during the first stages of their life cycle, prefer shallow waters where they can easily find preys and havens (Simpfendorfer & Milward 1993, Salomón-Aguilar *et al.* 2009).

A minimum size of 14.3 cm TL was recorded for *Z. xyster*. The smallest size recorded worldwide for this species was found in Costa Rican waters with 14.0 cm TL (Espinoza *et al.* 2015). Present results represent the smallest size in Ecuadorian waters, where there were records of a minimum size of 38 cm TL (Martínez-Ortiz & García-Domínguez 2013). Minimum and maximum total lengths of 21.0 and 118.0 cm, respectively, have been reported for *P. leucorhynchus* in Ecuadorian waters by Romero-Caicedo *et al.* (2015), which differs from reported in the current study, with a maximum total length of 76.1 cm, also reported in Colombian waters (76.0 cm LT) (Payán *et al.* 2011).

Length-weight relationships for the species showed  $r^2$  values ranging from 0.92 to 0.98, therefore, these estimations can be considered near to adequate. Analysis by sex also showed high values of  $r^2$  (0.97 to 0.99), except for males of *G. crebripunctata* ( $r^2 = 0.66$ ), which may be due to low sample number or the range of limited sizes for males. According to Froese *et al.* (2011), when determination coefficient ( $r^2$ ) is minor than 0.95 it may indicate outliers, due to the inclusion of extreme individuals such as juveniles (minor size) or adults (major size), abrupt change of shape during development, sex-differences, seasonal differences, among others.

First records of growth type are provided for *R. equatorialis* and *G. crebripunctata* (Table 1). Allometric growth parameter  $b$  for *R. equatorialis* was 3.11 (positive allometric growth) and *G. crebripunctata* showed isometric growth (2.94). For *Zapteryx xyster* values showed negative allometric growth ( $b = 2.92$ ), this type of growth is similar as the one reported in the Southwest Atlantic Ocean for *Z. brevirostris*, a congener species (Pasquino *et al.* 2016), but differs from the value reported in a previous work in the Colombian Pacific of  $b = 2.85$ , interpreted by the authors as isometric growth (Mejía-Falla *et al.* 2006). Isometric growth herein reported for *P. leucorhynchus* differs from other studies in Colombian and Ecuadorian Pacific, where allometric growth rates were reported by Payán *et al.* (2011) and Romero-Caicedo *et al.* (2015). Male and female presented the same type of growth, except for *G. crebripunctata*, where an isometric growth was observed for females ( $b = 3.12$ ), and negative allometric growth for males ( $b = 2.41$ ) (Table 1), however, as proposed by Froese *et al.* (2011), a re-evaluation of these type of cases is suggested.

**Table 1. Descriptive statistics and estimated parameters of length-weight relationships for four species of batoids collected from Ecuadorian Pacific Ocean / Estadística descriptiva y parámetros estimados de las relaciones longitud-peso para cuatro especies de batoideos recolectadas del océano Pacífico Ecuatoriano**

Species / Sex	n	Length (cm)		Weight (g)		Parameters							
		Min	Max	Min	Max	a	CI <sub>95%</sub> (a)	b	CI <sub>95%</sub> (b)	r <sup>2</sup>	t-test	P	Growth type
Gymnuridae													
<i>Gymnura crebripunctata</i> *	101	20.0	71.5	70.0	3665.0	0.0111	0.0058 - 0.0211	2.94	2.77 - 3.12	0.92	-0.65	>0.05	Isometric
Male*	49	20.0	46.0	70.0	900.0	0.0768	0.0123 - 0.4789	2.41	1.90 - 2.92	0.66	-2.34	<0.05	Allometric (-)
Female*	52	22.0	71.5	100.0	3665.0	0.0055	0.0030 - 0.0099	3.12	2.97 - 3.28	0.97	1.64	>0.05	Isometric
Rajidae													
<i>Rostroraja equatorialis</i>	519	7.8	61.0	4.6	1105.0	0.0036	0.0032 - 0.0041	3.11	3.08 - 3.15	0.98	6.61	<0.05	Allometric (+)
Male	230	10.1	58.2	5.2	1105.0	0.0033	0.0027 - 0.0040	3.15	3.10 - 3.20	0.98	5.53	<0.05	Allometric (+)
Female	289	7.8	61.0	4.6	1105.0	0.0039	0.0033 - 0.0045	3.09	3.05 - 3.14	0.99	4.13	<0.05	Allometric (+)
Rhinobatidae													
<i>Pseudobatos leucorhynchus</i>	297	22.3	76.1	30.0	1670.0	0.0029	0.0023 - 0.0035	3.05	3.00 - 3.10	0.97	1.78	>0.05	Isometric
Male	129	22.3	69.0	30.0	1550.0	0.0032	0.0022 - 0.0045	3.02	2.93 - 3.11	0.97	0.46	>0.05	Isometric
Female	168	23.1	76.1	30.0	1670.0	0.0027	0.0021 - 0.0035	3.06	3.00 - 3.13	0.98	1.94	>0.05	Isometric
Trygonorrhinidae													
<i>Zapteryx xyster</i>	507	14.3	66.0	17.3	1540.0	0.0079	0.0068 - 0.0092	2.92	2.88 - 2.96	0.98	-4.08	<0.05	Allometric (-)
Male	247	14.3	62.0	17.3	1205.0	0.0080	0.0062 - 0.0103	2.92	2.85 - 2.98	0.97	-2.47	<0.05	Allometric (-)
Female	260	16.0	66.0	24.5	1540.0	0.0079	0.0066 - 0.0095	2.92	2.87 - 2.96	0.98	-3.50	<0.05	Allometric (-)

n: sample size; a: intercept; b: slope; CI<sub>95%</sub>: confidence intervals; r<sup>2</sup>: coefficient of determination; Student-t: test to evaluate growth  
P: significance value of the Student-t test and growth type; Species/Sex\*: disc width

According to ANCOVA analyse, significant differences were not found in LWR between sex for three species (*P. leucorhynchus*:  $F= 0.055$ ,  $P > 0.05$ ; *Z. xyster*:  $F= 2.498$ ,  $P > 0.05$ ; *R. equatorialis*:  $F= 0.088$ ,  $P > 0.05$ ), while *G. crebripunctata* did present differences ( $F= 9.706$   $P < 0.05$ ), in which males had a negative allometric growth. Our results with other studies show that the values of *b* can be affected by several factors, such as differences in the number of examined organisms, area, season, or the use of different lengths (Froese *et al.* 2011, Pan *et al.* 2014, Ehemann *et al.* 2017).

This study represents the first contribution to knowledge of biological aspects for *G. crebripunctata* and *R. equatorialis*; as well as an increase in the life history information of *P. leucorhynchus* and *Z. xyster* in the Ecuadorian waters. The data obtained constitutes a baseline for future models used for fisheries resources management and conservation, however, it is suggested that the results of these relationships be used only with the presented length ranges.

## ACKNOWLEDGMENTS

A special thanks to the crew of the boats that allowed us to collect the biological information of the species used in this work, as well to Robert Leaf, Víctor Cruz Escalona, Francisco Arreguín Sánchez and Olivia Echazabal Salazar for their help in translating this article to English and for their valuable comments that allowed to enrich and improve the manuscript.

## LITERATURE CITED

- Correia E, JP Granadeiro, A Regalla & P Catry. 2018. Length-weight relationship of fish species from the Bijagós Archipelago, Guinea-Bissau. *Journal of Applied Ichthyology* 34(1): 177-179.
- Dulvy NK, SL Fowler, JA Musick, RD Cavanagh, PM Kyne, LR Harrison, JK Carlson, LNK Davidson, SV Fordham, MP Francis, CM Pollock, CA Simpfendorfer, GH Burgess, KE Carpenter, LJV Compagno, DA Ebert, C Gibson, MR Heupel, SR Livingstone, JC Sanciangco, JD Stevens, S Valenti & WT White. 2014. Extinction risk and conservation of the world's sharks and rays. *eLife* 3: e00590. <<https://doi.org/10.7554/eLife.00590.001>>

- Ehemann NR, XA Pérez-Palafox, P Mora-Zamacona, MI Burgos-Vázquez, AF Navia, PA Mejía-Falla & VH Cruz-Escalona. 2017.** Size-weight relationships of batoids captured by artisanal fishery in the southern Gulf of California, Mexico. *Journal of Applied Ichthyology* 33(5): 1051-1054.
- Espinoza M, SEM Munroe, TM Clarke, AT Fisk & IS Wehrmann. 2015.** Feeding ecology of common demersal elasmobranch species in the Pacific coast of Costa Rica inferred from stable isotope and stomach content analyses. *Journal of Experimental Marine Biology and Ecology* 470: 12-25.
- Froese R. 2006.** Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology* 22(4): 241-253.
- Froese R, AC Tsikliras & KI Stergiou. 2011.** Editorial note on weight-length relations of fishes. *Acta Ichthyologica Et Piscatoria* 41: 261-263.
- Gladston Y, KV Akhilesh, C Thakurdas, OPK Ravi, SMAjina & L Shenoy. 2018.** Length-weight relationship of selected elasmobranch species from north-eastern Arabian Sea, India. *Journal of Applied Ichthyology* 34(3): 753-757.
- González-González LDV & NREhemann. 2019.** Length-weight relationships of six elasmobranch species captured at the artisanal fishery of Margarita Island, Venezuela. *Journal of Applied Ichthyology* 35(2): 594-596.
- Koushlesh SK, A Sinha, K Kumari, S Borah, TN Chanu, R Baitha, SK Das, P Gogoi, SK Sharma, MH Ramteke & BK Das. 2017.** Length-weight relationship and relative condition factor of five indigenous fish species from Torsa River, West Bengal, India. *Journal of Applied Ichthyology* 34(1): 169-171.
- Last P, G Naylor, B Séret, W White, M de Carvalho & M Stehmann. 2016.** *Rays of the World*, 1577 pp. CSIRO Publishing, Melbourne.
- Liu C, Y Wang, Q Ren, W Li, Z Jiang & W Xian. 2018.** Length-weight relationships of three freshwater fish species from the Songhua River and Nen River, China. *Journal of Applied Ichthyology* 34(3): 737-738.
- Martínez-Ortiz J & M García-Domínguez. 2013.** *Chondrichthyes of Ecuador Field Guide. Chimaeras, Sharks and Rays*, 246 pp. Ministry of Agriculture, Livestock, Aquaculture and Fisheries (MAGAP), Vice Ministry of Aquaculture and Fisheries (VMAP), Undersecretariat of Fisheries Resources (SRP), Manta.
- Mejía-Falla PA, AF Navia & A Giraldo. 2006.** Notas biológicas de la raya ocelada *Zapteryx xyster* Jordan & Evermann, 1896 (Chondrichthyes: Rhinobatidae) en la zona central de pesca del Pacífico colombiano. *Investigaciones Marinas* 34(2): 181-185.
- Ogle DH, P Wheeler & A Dinno. 2018.** *FSA: Fisheries Stock Analysis*. R package version 0.8.22. <<https://github.com/droglenc/FSA>>
- Pan L, JJ Xie, Z Yang, HY Tang & Y Qiao. 2014.** Length-weight relationships of six fish species from the upper reaches of the Yangtze River, southwest China. *Journal of Applied Ichthyology* 30(3): 552-554.
- Pasquino AF, MF Martins & OBF Gadig. 2016.** Length-weight relationship of *Rhinobatos horkelii* Müller & Henle, 1841 and *Zapteryx brevirostris* (Müller & Henle, 1841) off Brazil, southwestern Atlantic Ocean. *Journal of Applied Ichthyology* 32(6): 1282-1283.
- Payán LF, AF Navia, EA Rubio & PA Mejía-Falla. 2011.** Biología de la raya guitarra *Rhinobatos leucorhynchus* (Günther, 1867) (Rajiformes: Rhinobatidae) en el Pacífico colombiano. *Latin American Journal of Aquatic Research* 39(2): 286-296.
- R Core Team. 2013.** R: a language and environment for statistical computing. R Foundation for Statistical Computing, Viena. <<http://www.R-project.org>>
- Rastgoo AR, MR Fatemi, T Valinassab & MS Mortazavi. 2016.** Length-weight relationships for 10 elasmobranch species from the Oman Sea. *Journal of Applied Ichthyology* 32(4): 734-736.
- Romero-Caicedo AF, P Looor-Andrade, A Cruz-Martínez & M Carrera-Fernández. 2015.** Weight-length relationships of six batoids in the Ecuadorian Pacific. *Journal of Applied Ichthyology* 31(5): 965-966.
- Salomón-Aguilar CA, CJ Villavicencio-Garayzar & H Reyes-Bonilla. 2009.** Shark breeding grounds and seasons in the Gulf of California: Fishery management and conservation strategy. *Ciencias Marinas* 35(4): 369-388.
- Simpfendorfer CA & NE Milward. 1993.** Utilisation of a tropical bay as a nursery area by sharks of the families Carcharhinidae and Sphyrnidae. *Environmental Biology of Fishes* 37(4): 337-345.
- Smith WD, JJ Bizzarro, VP Richards, J Nielsen, F Márquez-Farías & MS Shivji. 2009.** Morphometric convergence and molecular divergence: the taxonomic status and evolutionary history of *Gymnura crebripunctata* and *Gymnura marmorata* in the Eastern Pacific Ocean. *Journal of Fish Biology* 75(4): 761-783.
- Wang PG, DJ Booker, SK Crow & DJ Jellyman. 2013.** Does one size fit all? An evaluation of length-weight relationships for New Zealand's freshwater fish species. *New Zealand Journal of Marine and Freshwater Research* 47(4): 450-468.
- Wei N, Y Zhang, F Wu, Z Shen, H Ru, L Liu & Z Ni. 2019.** Length-weight relationships for 15 fish species from the Three Gorges Reservoir. *Journal of Applied Ichthyology* 35(3): 789-792.
- Zar JH. 1999.** *Biostatistical analysis*, 663 pp. Prentice Hall, Upper Saddle River.

---

Received 9 March 2020

Accepted 27 September 2021